



## 2002 INDIANA ADULT TOBACCO SURVEY

The Indiana Adult Tobacco Survey (ATS) was conducted from October 30 to December 22, 2002 surveying 1903 Indiana adults, including an over sample of African Americans and Hispanics. ATS, developed by the Centers for Disease Control and Prevention (CDC), was adapted by ITPC by adding questions designed specifically for Indiana in order to serve as a surveillance measure for statewide tobacco use prevalence among adults.

### County breakouts for ATS geographic regions<sup>1</sup>

- **Northwest: 33.0%**
  - Lake, Porter, Newton, Jasper, LaPorte
- **North Central: 23.8%**
  - St. Joseph, Elkhart, Starke, Marshall, Kosciusko, Pulaski, Fulton, White, Carroll, Cass, Miami, Wabash
- **Northeast: 21.3%**
  - LaGrange, Steuben, Noble, DeKalb, Whitley, Allen, Huntington, Wells, Adams
- **Central West: 32.0%**
  - Benton, Warren, Tippecanoe, Fountain, Montgomery, Vermillion, Parke, Putnam, Vigo, Clay, Owen, Sullivan, Greene
- **Central-Indianapolis Metropolitan Area: 26.6%**
  - Marion, Boone, Hamilton, Madison, Hancock, Shelby, Johnson, Morgan, Hendricks, Clinton, Howard, Tipton
- **Central East: 30.4%**
  - Grant, Blackford, Jay, Delaware, Randolph, Henry, Wayne, Rush, Fayette, Union, Franklin
- **Southwest: 25.6%**
  - Monroe, Brown, Lawrence, Knox, Daviess, Martin, Orange, Gibson, Pike, Dubois, Crawford, Posey, Vanderburgh, Warrick, Spencer, Perry
- **Southeast: 33.0%**
  - Bartholomew, Decatur, Jackson, Jennings, Ripley, Dearborn, Washington, Scott, Jefferson, Switzerland, Ohio, Harrison, Floyd, Clark

---

<sup>1</sup> The differences between geographic regions are not statistically different from each other or to the overall state current smoking prevalence.